
Quantum magnetism in two dimensions: From semi-classical Néel order to magnetic disorder

J. Richter¹, J. Schulenburg² and A. Honecker³

¹ Institut für Theoretische Physik, Otto-von-Guericke-Universität Magdeburg,
P.O.Box 4120, D-39016 Magdeburg, Germany
johannes.richter@physik.uni-magdeburg.de

² Universitätsrechenzentrum, Otto-von-Guericke-Universität Magdeburg, P.O.Box
4120, D-39016 Magdeburg, Germany
joerg.schulenburg@urz.uni-magdeburg.de

³ Institut für Theoretische Physik, TU Braunschweig, Mendelssohnstr. 3, D-38106
Braunschweig, Germany & Universität Hannover, Institut für Theoretische
Physik, Appelstrasse 2, D-30167 Hannover, Germany
a.honecker@tu-bs.de

Published version: Chapter 2 in *Quantum Magnetism*, ed by U. Schollwöck,
J. Richter, D.J.J. Farnell, R.F. Bishop, Lecture notes in physics **645**, 85-153
(Springer-Verlag, Berlin Heidelberg 2004)

Abstract:

In this article we focus on the ground state and the low-lying excitations of the $s = 1/2$ Heisenberg antiferromagnet (HAFM) on the 11 two-dimensional (2D) uniform Archimedean lattices.

Although we know from the Mermin-Wagner theorem that thermal fluctuations are strong enough to destroy magnetic long-range order (LRO) for Heisenberg spin systems at any finite temperature in one and two dimensions, the role of quantum fluctuations is less understood. While the ground state of the one-dimensional (1D) quantum HAFM is not long-range ordered, the quantum HAFM e.g. on the 2D square and triangular lattices exhibits semi-classical Néel like LRO. However, in two dimensions there are many other lattices with different coordination numbers and topologies, and there is no general statement concerning zero-temperature Néel-like LRO. Recent experimental results on CaV_4O_9 and $\text{SrCu}_2(\text{BO}_3)_2$ demonstrate the possibility of non-Néel ordered ground states and signal that the $s = 1/2$ HAFM on 2D lattices with appropriate topology may have a ground state without semi-classical LRO.

Based on extensive large-scale exact diagonalization studies of the ground state and the low-lying excitations for the spin-1/2 HAFM on the Archimedean

lattices we compare and discuss the ground-state features of all 11 lattices. In this manner we obtain some insight in the influence of lattice topology on magnetic ordering of quantum antiferromagnets in two dimensions. From our results we conclude that the ground state of the spin-1/2 HAFM on most of the Archimedean lattices (in particular the four bipartite ones) turns out to be semi-classically Néel-like ordered. However, we find that the interplay of competition of bonds (geometric frustration and non-equivalent nearest neighbor bonds) and quantum fluctuations gives rise to a quantum paramagnetic ground state without semi-classical LRO for two lattices. The first one is the famous kagomé lattice, for which this statement is well-known by numerous studies during the last decade. Remarkably, we find one additional lattice among the 11 uniform Archimedean lattices, the so-called star lattice, with a quantum paramagnetic ground state. For both these Archimedean lattices the ground state is highly degenerate in the classical limit $s \rightarrow \infty$, although notably their quantum ground states are fundamentally different.

Furthermore, we present numerical results for the magnetization curve of the HAFM on all 11 Archimedean lattices. The magnetization process is discussed in some detail for the square, triangular and kagomé lattices. One focus are plateaus appearing in the magnetization curve due to quantum fluctuations and geometric frustration. In particular, the kagomé lattice exhibits a rich spectrum of magnetization plateaus. Another focus are magnetization jumps arising on the kagomé and the star lattice just below the saturation field. These magnetization jumps may be understood analytically by using independent local magnon excitations.

Some related $s = 1/2$ models are also discussed briefly. Particular attention is given to the 2D Shastry-Sutherland model. For this model, we discuss quantum phase transitions and discuss the magnetization curve in comparison with experiments on $\text{SrCu}_2(\text{BO}_3)_2$.

Contents (preprint version)

1	Introduction	4
2	Archimedean Lattices	6
2.1	Characteristics and geometry	6
2.2	Relationships between the lattices	8
3	Criteria for Néel like order	11
3.1	Order parameter	11
3.2	Mechanism of symmetry breaking - the Pisa tower of quasi-degenerate joint states (QDJS)	13
3.3	Finite-size scaling	16
4	Magnetic ground-state ordering for the spin half HAFM on the Archimedean lattices	18
4.1	Semi-classical Néel ordering on bipartite lattices	18
4.2	Semi-classical LRO on frustrated lattices	25
4.3	Absence of semi-classical LRO on frustrated lattices - the kagomé (T8) and the star (T9) lattices	34
4.4	Summary and comparison	39
5	Quantum phase transitions in 2D HAFM - the CaVO $J - J'$ model and the Shastry-Sutherland model	44
6	Magnetization process	48
6.1	Square lattice	50
6.2	Triangular lattice	53
6.3	Kagomé lattice	55
6.4	Independent magnons and macroscopic magnetization jumps ..	57
6.5	Shastry-Sutherland model versus $\text{SrCu}_2(\text{BO}_3)_2$	61
6.6	Summary of plateaux & related topics	64
	References	66

Complete preprint version of this article:

PostScript at: <http://www.tu-bs.de/~honecker/papers/2dqm.ps.gz>
 PDF at: <http://www.tu-bs.de/~honecker/papers/2dqm.pdf>

Complete published version of this article available from:

<http://www.springerlink.com/link.asp?id=ccv22wcn1jnrxfhu>
 PDF of published version also at:
<http://www.tu-bs.de/~honecker/papers/lmp645.pdf>

Additional features of preprint version:

- Extended Abstract,
- References updated,
- Separate Table of Contents,
- Value of the square of the order parameter $N = 36$ kagomé lattice corrected: $(m^+)^2 = 0.059128$.